

In recent years other *Cupressus* feeding moths have been recorded on Guernsey, *Eupithecia phoeniceata* was first recorded in 1959 and is now common, and *Lithophane leautieri hesperica* which was first noted in 1971 is similarly frequently seen. Whether or not Channel Island insects are British and therefore whether *T. cupressata* can now be added to the list of British moths is a matter for discussion but we noted with interest that the recent discovery of *Eupithecia ultimaria* on Guernsey was described as an addition to the British list.

### Acknowledgements

We would like to thank the staff of the British Museum (Natural History) for identifying the specimens, and Mr. Barry Goater for confirming the identification and for providing help with the references. We also thank Mr. Graham Jackson for providing the photograph.

### References

- Culot, J., 1917-1919. Noctuelles et Géomètres d'Europe I: 20 pl. 23 fig. 465.  
Seitz, A., 1912. The Macrolepidoptera of the World 4: 218 pl. 8(i).  
Peet, T. N. D., 1986. An early record of *Eupithecia phoeniceata* (Rambur) (Lepidoptera: Geometridae) in Guernsey. *Entomologist's Gaz.* 37: 60.  
Peet, T. N. D., In press. *Rep. Soc. guernes.* Entomology Report for 1985.  
Riley, A. M., 1985. *Eupithecia ultimaria* Boisduval (Lepidoptera: Geometridae): A pug new to the British list. *Entomologist's Gaz.* 36: 259.

**Note:** A further two specimens of *cupressata* were taken at the same site on the nights of 10th and 11th July 1986. P.D.M.C.

---

DAMAGE TO BLUEBERRY (*VACCINIUM CORYMBOSUM*) BY *CACOECIMORHA PRONUBANA* (HÜBNER) — On 20th May, 1986 tortricid larvae was found on various cultivars of highbush blueberry in a plantation at East Malling Research Station, where they had caused extensive damage to young shoots, leaves and flowers. Most of the larvae were almost fully-grown and most had constructed shelters from dead flowers or leaves webbed to the shoots. Ten larvae were collected and reared through to adult; all proved to be *Cacoecimorpha pronubana* (Hubner), which is known to be polyphagous. M. A. EASTERBROOKE, East Malling Research Station, nr. Maidstone, Kent.